



Health Record #: _____	Complete or place barcoded patient label here	
Patient Name: <i>(Print first, last)</i> _____		
DOB: <u>dd</u> / <u>mm</u> / <u>yy</u>	Age: _____	<input type="checkbox"/> Female <input type="checkbox"/> Male
OHIP #: _____	Version Code: _____	
Account #: _____	Date of Admission: <u>dd</u> / <u>mm</u> / <u>yy</u>	

Electroencephalogram (EEG) Requisition

Previous EEG? <input type="checkbox"/> Yes <input type="checkbox"/> No If Yes, provide Date: <u>dd</u> / <u>mm</u> / <u>yy</u>	
To book an appointment call (905) 895-4521, ext. 2665. Fax completed requisitions to (905) 830-5965	
Provisional Diagnosis:	
History:	
Medications:	
Positive Findings:	
Physician Name: <i>(print first, last)</i>	
Signature:	Date: <u>dd</u> / <u>mm</u> / <u>yy</u> Time: _____

ATTENTION PHYSICIAN

1. Preparation for EEG

- a) Please ask your patient to eat a regular meal (or at least a snack) before the EEG appointment time to help stabilize blood sugar level.
- b) The patient must wash their hair thoroughly before the test.
- c) Patients should continue to take their normal medications, unless directed otherwise.

2. For child sleep-deprived EEG, please instruct parent(s) to follow instructions below: **Book Appointment Times**

<i>newborn - 3 years</i>	<i>Do not let nap in the a.m.</i>	<i>Afternoon</i>
<i>4 - 7 years</i>	<i>Keep child awake from 2 a.m.</i>	<i>Morning</i>
<i>8 - 12 years</i>	<i>Keep child awake from midnight</i>	<i>Morning</i>
<i>greater than 12 years</i>	<i>Keep child awake all night</i>	<i>Morning</i>

**Do not give child any food or drink containing caffeine (e.g. coke, chocolate, tea, coffee) prior to test. Do not let child sleep on way to hospital.*

- ### 3. In rare circumstances, sedation for paediatric patients may be required.
- If so, please call the patient scheduling department at extension 2665 to make the necessary special arrangements for nursing availability during the procedure. The drug of choice for EEG is chloral hydrate, suggestive dosage 25 to 50 mg per kg, available as syrup 250mg, 500 mg and capsules 500mg.